

LA-UR-17-21138

Approved for public release; distribution is unlimited.

Title: The Scientific Challenges in Stewarding the U.S. Nuclear Weapons

Stockpile

Author(s): Wilburn, Wesley Scott

Intended for: Colloquium at Indiana University

Issued: 2017-02-14



The Scientific Challenges in Stewarding the U.S. Nuclear Weapons Stockpile

W.S. Wilburn

Deputy Division Leader for Predictive Capability

X Theoretical Division
Los Alamos National Laboratory





LANL Stewards Much of the U.S. Nuclear Stockpile

- We are the design agency for four of seven weapons systems in the national arsenal
- We are responsible for the safety, surety, and reliability of these systems



W78 land-based warhead



B61 aircraft-carried bomb



W76/W88 sub-based warheads

Since 1992, we have performed our mission without nuclear testing




Nuclear Weapons Release Nuclear Binding Energy


- Both heavy-nuclei fission and light-nuclei fusion release energy
- ullet Fission releases \sim 200 MeV per fission event
 - Example: $n+^{235}U \rightarrow^{236}U \rightarrow^{92}Kr +^{141}Ba +3n$
 - Note that additional neutrons are produced, inducing further fissions
- Fusion produces 17.6 MeV per fusion (for D+T)
 - Example: D + T \rightarrow 4He +n
- Compare to chemical reactions that release eV's per reaction event



Early Nuclear Weapons Used Only Fission

- Function by creating a super-critical mass (neutron leakage rate < production rate)
- Two technologies were developed by LANL during the Manhattan Project
- Gun weapon assembles sub-critical pieces of fissile material into super-critical mass
- Implosion weapon compresses sub-critical piece of fissile material into super-critical mass








Little Boy Was the First Gun-Assembled Weapon


- Simple design (no proof test required)
- Limited to ²³⁵U to avoid pre-initiation
- Inefficient use of nuclear material
- Hiroshima August 6, 1945
- Yield of 15 kt






Trinity/Fat Man Was the First Gun-Assembled Weapon

- Complex design using precision explosives
- Can use ²³⁵U or ²³⁹Pu
- Efficient use of nuclear material
- Trinity July 16, 1945
- Nagasaki August 9, 1945
- Yield of 21 kt





Modern Weapons are Thermonuclear




- Thermonuclear weapons have two physically separate stages: primary and secondary
- The primary fission stage goes off first, followed by the secondary stage
- X-ray energy produced by the primary stage heats and compresses the secondary stage
- The device produces energy by fission and fusion
- Much higher yields possible



Ivy/Mike Was the First Thermonuclear Test

- Experimental device
- Used liquid D₂ as TN fuel
- Not suitable as a weapon
- Enewetok October 31, 1952
- Yield of 10.4 Mt







Castle/Bravo Was the Largest U.S. Nuclear Test

- Bikini February 28, 1954
- Predicted yield of 6 MT
- Actual yield of 15 Mt






World Record for Largest Nuclear Test Held by USSR

- "Tsar Bomba"
- Novaja Zemlya October 30, 1961
- Design Yield 100 Mt
- Tested at reduced yield of 58 Mt





Atmospheric testing was banned by treaty in 1963




Current U.S. Stockpile Was Designed for Cold War

- Designed for maximum yield in minimum size and weight
 - Highly optimized
- Intended to be replaced after 15-20 years
 - Now well past original design life
- Use hazardous materials
 - Expensive and difficult to handle today
- Very complex, with many parts
 - Challenging to maintain without nuclear testing



B61 Nuclear Bomb has More Than 4000 Parts





Why is Maintaining the Stockpile So Hard?

- Operating conditions of a nuclear weapon exist nowhere else and cannot be fully replicated in a lab setting
 - Temperatures $> 10^8$ K
 - Material velocities > 10⁶ m/s
 - Pressures $> 10^7$ bar
 - Time scales $< 10^{-8}$ s
- We have developed a science-based method to stewarding the stockpile
 - Use large-scale multi-physics simulations to predict weapon performance
 - Perform small-scale experiments to continuously improve our understanding of the relevant physics
 - Validate the simulations against legacy test data and integrated non-nuclear and sub-critical experiments
- We also have a surveillance to monitor for unexpected changes



Computer Simulations of Nuclear Weapons


- We use computational models
 - High explosive burn
 - Radiation/Hydrodynamics
 - Neutronics
 - Thermonuclear burn
- Incorporating physical data
 - Neutron cross sections
 - Equation of state
 - Opacities
- Running on high-performance computers
 - World's first computer > 1 Petaflop
 - LANL is part of the Exascale Computing Project







We Compare Our Simulations to Legacy Test Data










1054 nuclear tests (24 joint with UK)



Dual-Axis Radiographic Hydrodynamic Test Facility





- DARHT provides two beams of high-energy pulsed x-rays
- Allows testing the dynamics of implosions using non-nuclear components as a function of time
- These experiments are a critical part of validating our simulations



Sub-Critical Experiments Study Nuclear Materials





- Conducted underground at the Nevada National Security Site (formerly NTS)
 - The experiments are performed at a depth of 300 m to contain any radioactive release
- We can test the initial stages of implosion using plutonium
 - The amount of plutonium is reduced to avoid criticality
- Dynamic plutonium experiments provide important information on this material



High-Energy X-Rays and Laser Probes Provide Data





- Measure properties such as
 - Equations-of-state
 - Cavity formation and strength
 - Spall and damage
 - Ejecta
 - Material properties of aged versus new plutonium
 - Material properties of cast versus wrought plutonium
 - Metallurgy and its effects on other dynamic properties



Example Radiographic Images




- Experiment designed to test reproducibility of dynamic experiments
- Two identical packages with non-nuclear materials were shot simultaneously
- High explosives drove shock through material from the bottom



Multi-Point Photon Doppler Velocimetry





- Multiple laser beams allow velocity measurements of material at multiple points
- Doppler shift of beam reflecting from moving surface gives velocity
- Provides a rich data set (3M velocities) to compare to simulations




Small-Scale Experiments









Experiments improve the science behind nuclear weapons




Proton Radiography Studies Dynamic Materials



- pRad uses 800 MeV proton beam to image dynamic experiments
- ullet Beam can be pulsed to provide 40 images \geq 100ns apart
- Images are analyzed to provide density information as a function of position and time




pRad Uses Multiple Coulomb Scattering






pRad Example: Melting and Solidification of Al-In Alloy






pRad Example: Instability Growth




• Shock applied to two materials with different densities



Precision Measurements of Nuclear Cross Sections






- The 239 Pu(n, f) cross section affects nuclear weapons performance
- Spread in existing data sets suggests uncontrolled or unaccounted for systematic uncertainties
- Lawrence Livermore National Laboratory and LANL are measuring this cross section at LANSCE to 1% precision




Experimental Facilities to Study Radiation Flow




Z-Pinch
Sandia National Laboratory
Albuquerque, New Mexico





National Ignition Facility
Lawrence Livermore National Laboratory
Livermore, California





Summary

- Maintaining the U.S. nuclear stockpile without nuclear testing is very challenging
- We use a variety of science-based tools to perform this mission
- Large-scale multi-physics simulations running on high-performance computers have replaced nuclear testing
- Experimental data is crucial to calibrate and validate the simulations
 - Data from legacy nuclear tests
 - Large-scale integrated experiments
 - Sub-critical experiments
 - Small-scale experiments
- Many scientific disciplines contribute to maintaining the stockpile

